

| Drug                   | Treatment For                           | Dosage Form          | Covered as Rx | Self-Injectable Product | Costs \$600 or more for up to a 30 day supply | Special Handling (Refrigerate, Limited Distribution, etc)   | Limited Distribution by Manufacturer | Available from Wholesaler |
|------------------------|---|----------------------|---------------|-------------------------|---|---|--------------------------------------|---------------------------|
| Abraxane               | Oncology- Injectable                    | IV                   | No            | No                      | Yes   | Store at room temp. Protect from bright light. Freezing or refrigerating do not adversely affect the stability of intact vials. Different standards apply after reconstitution  | No                                   | Yes                       |
| Adagen                 | Immune Deficiencies & Related Disorders | Injection IM         | No            | No                      | Yes   | Refrigerate; do Not freeze. Discard unused portions; do Not save for further use. Do Not use if previously frozen   | No                                   | No                        |
| Adcirca                | Pulmonary Arterial Hypertension         | Oral                 | Yes           | No                      | Yes   |   | No                                   | Yes                       |
| Afinitor               | Oncology- Oral/Topical                  | Oral                 | Yes           | No                      | Yes   | Protect from light; protect from moisture. Reconstituted solution and infusion solution are stable for 24 hours when refrigerated at 2°C to 8°C   | No                                   | Yes                       |
| Alimta                 | Oncology- Injectable                    | IV                   | No            | No                      | Yes   | none  | No                                   | Yes                       |
| Alunbrig               | Oncology- Oral/Topical                  | Oral                 | Yes           | No                      | Yes   | Store at 2°C to 8°C (36°F to 46°F); do Not freeze. Do Not shake. Protect from light. Store in original carton until use. The following stability information has also been reported: May be stored at room temperature for up to 7 days | No                                   | No                        |
| Aranesp                | Anemia                                  | Injection SubQ or IV | Yes           | No                      | Yes   | Refrigerate, Do not freeze. Protect from light. Vial: If refrigeration is not available, may be stored at 25°C (77°F) for up to 30 days   | No                                   | Yes                       |
| Avonex                 | Multiple Sclerosis                      | Injection IM         | Yes           | Yes                     | Yes   | Prior to reconstitution, store intact vials at room temperature of 25°C; excursions permitted to 15°C to 30°C   | No                                   | No                        |
| Azacitidine (Vidaza)   | Oncology- Injectable                    | Injection SubQ or IV | Yes           | Yes                     | Yes   | Refrigerate. Protect from light. Dispense in original container if applicable   | No                                   | No                        |
| Benlysta               | Systemic Lupus Erythematosus            | Injection SubQ or IV | No            | No                      | Yes   | If not used immediately following reconstitution, refrigerate solution at 2°C to 8°C and use within 3 hours; do not freeze or shake solution.   | No                                   | Yes                       |
| Betaseron              | Multiple Sclerosis                      | Injection SubQ       | Yes           | Yes                     | No  | Store at 2°C to 25°C (36°F to 77°F). Protect from light. Avoid humidity and high temperatures after opening bottle  | No                                   | Yes                       |
| Bexarotene (Targretin) | Oncology- Oral/Topical                  | Oral                 | Yes           | No                      | Yes   | none  | No                                   | Yes                       |
| Bosulif                | Oncology- Oral/Topical                  | Oral                 | Yes           | No                      | Yes   |   | No                                   | No                        |

| Drug                                      | Treatment For                                     | Dosage Form                  | Covered as Rx | Self-Injectable Product | Costs \$600 or more for up to a 30 day supply | Special Handling (Refrigerate, Limited Distribution, etc)  | Limited Distribution by Manufacturer | Available from Wholesaler |
|---|---|------------------------------|---------------|-------------------------|---|--|--------------------------------------|---------------------------|
| Botox                                     | Batten Disease                                    | Injection IM                 | No            | No                      | No  | Store unopened vials under refrigeration at 2°C to 8°C (36°F to 46°F) for up to 36 months or until the expiration date on the vial. After reconstitution, store in refrigerator at 2°C to 8°C (36°F to 46°F) and use within 24 hours (does Not contain a preservative).<br>Botox 100-unit vial: The following stability information has also been reported: May be stored at room temperature for up to 5 day<br>Lyophilized powder may be stored in the refrigerator or at room temperature of 3°C to 25°C. Protect from light. Discard unused portion. | No                                   | Yes                       |
| Bravelle<br>Capecitabine (Xeloda)         | Infertility<br>Oncology- Oral/Topical             | Injection SubQ or IM<br>Oral | Yes<br>Yes    | Yes<br>No               | No<br>Yes                                     | Prior to reconstitution, store at or below 30°C (86°F). Do not freeze. Following reconstitution in a sterile laminar air flow environment, store under refrigeration. Don't freeze, shake, or heat<br>Lyophilized powder may be stored in the refrigerator or at room temperature of 3°C to 25°C. Protect from light. Discard unused portion.  | No<br>No                             | Yes<br>Yes                |
| Carimune NF                               | Immune Deficiencies & Related Disorders           | IV                           | No            | No                      | Yes   | Store intact vials and syringes at 2°C to 8°C; do Not freeze. Do Not separate contents of carton prior to use. Protect from light. Bring to room temperature prior to administration. Store reconstituted solution for up to 24 hours at 2°C to 8°C (do Not freeze); Do Not leave at room temperature for more than 2 hours prior to administration.   | No                                   | Yes                       |
| Cetrotide                                 | Infertility                                       | Injection SubQ               | Yes           | Yes                     | No  | Store at intact vials at 15°C to 30°C. Following reconstitution, solution is stable when refrigerated for 30 days (Novarel) or 60 days (Pregnyl).  | No                                   | Yes                       |
| Chorionic gonadotropin (Novarel, Pregnyl) | Infertility                                       | Injection IM                 | Yes           | Yes                     | No  | Store intact vials and syringes at 2°C to 8°C; do Not freeze. Do Not separate contents of carton prior to use. Protect from light. Bring to room temperature prior to administration. Store reconstituted solution for up to 24 hours at 2°C to 8°C (do Not freeze); Do Not leave at room temperature for more than 2 hours prior to administration.   | No                                   | Yes                       |
| Cimzia                                    | Inflammatory Bowel Disease / Rheumatoid Arthritis | Injection SubQ               | Yes           | Yes                     | Yes   | Store at 2°C to 8°C (36°F to 46°F) for up to 36 months or at room temperature for up to 12 months; do Not return vial to refrigerator after it has been stored at room temperature. Do Not freeze. Keep in original carton to protect from light.  | No                                   | Yes                       |
| Cuvitru                                   | Transplant  | Injection SubQ or IM         | No            | No                      | Yes   | Store at 2°C to 8°C (36°F to 46°F) for up to 36 months or at room temperature for up to 12 months; do Not return vial to refrigerator after it has been stored at room temperature. Do Not freeze. Keep in original carton to protect from light.  | No                                   | No                        |
| Daklinza                                  | Hepatitis C                                       | Oral                         | Yes           | No                      | Yes   | None   | No                                   | Yes                       |
| Dofetilide (Tikosyn)                      | Cardiac Disorders                                 | Oral                         | Yes           | No                      | No  | None   | No                                   | Yes                       |

| Drug                                      | Treatment For                    | Dosage Form               | Covered as Rx | Self-Injectable Product | Costs \$600 or more for up to a 30 day supply | Special Handling (Refrigerate, Limited Distribution, etc)   | Limited Distribution by Manufacturer | Available from Wholesaler |
|---|----------------------------------|---------------------------|---------------|-------------------------|---|---|--------------------------------------|---------------------------|
| Dupixent                                  | Atopic Dermatitis                | Injection SubQ            | Yes           | Yes                     | Yes   | N/A   | No                                   | No                        |
| Dysport                                   | Batten Disease                   | Injection IM              | No            | No                      | Yes   | Refrigerate, protect from light   | No                                   | No                        |
| Eligard (Canadian Brand name, See Lupron) | Hormonal Therapies               | Injection SubQ            | No            | No                      | Yes   | Refrigerate at 2°C to 8°C. Do Not shake. Do Not freeze or store in extreme heat or cold. Store in the original carton to protect from light or physical damage.   | No                                   | No                        |
| Enbrel                                    | Psoriasis / Rheumatoid Arthritis | Injection SubQ            | Yes           | Yes                     | Yes   | Individual autoinjectors, prefilled syringes, dose trays (containing multi-use vials and diluent syringes), or prefilled pens may be stored between 20°C and 25°C (68°F and 77°F) for a maximum single period of 14 days (Enbrel) with protection from light and sources of heat and humidity. Once an autoinjector, prefilled syringe, dose tray, or prefilled pen has been stored at room temperature, it should Not be placed back into the refrigerator; discard after 14 days (Enbrel) | No                                   | Yes                       |
| Entyvio                                   | Inflammatory Bowel Disease       | IV                        | No            | No                      | Yes   | Refrigerate unopened vials. Retain in original package to protect from light. After reconstitution and dilution, the infusion solution may be refrigerated for up to 4 hours). Don't freeze. Discard any unused portion   | No                                   | No                        |
| Epclusa                                   | Hepatitis C                      | Oral                      | Yes           | No                      | Yes   | Dispense in original container  | No                                   | Yes                       |
| Epogen                                    | Anemia                           | Injection SubQ or IV      | Yes           | Yes                     | Yes   | Vials should be stored at 2°C to 8°C (36°F to 46°F). Do Not freeze. Do Not shake. Protect from light.   | No                                   | Yes                       |
| Erwinaze                                  | Oncology- Injectable             | Injection IM or IV        | No            | No                      | Yes   | Refrigerate. Protect from light. Within 15 minutes of reconstitution, withdraw appropriate volume for dose into a polypropylene syringe. Do not freeze or refrigerate reconstituted solution; discard if not administered within 4 hours.   | No                                   | No                        |
| Euflexxa                                  | Osteoarthritis                   | Injection Intra-articular | No            | No                      | No  | Store refrigerated or at room temperature; do not freeze. Protect from light. If refrigerated, remove from refrigeration at least 20 to 30 minutes before use.  | No                                   | Yes                       |

| Drug           | Treatment For                           | Dosage Form               | Covered as Rx | Self-Injectable Product | Costs \$600 or more for up to a 30 day supply | Special Handling (Refrigerate, Limited Distribution, etc)  | Limited Distribution by Manufacturer | Available from Wholesaler |               |
|----------------|---|---------------------------|---------------|-------------------------|---|--|--------------------------------------|---------------------------|---------------|
| Exondys        | Duchenne Muscular Dystrophy             | IV                        | No            | No                      | Yes   | Store at 2°C to 8°C. Do Not freeze. Protect from light and store in the original carton until ready for use. The diluted solution may be stored at 2°C to 8°C for up to 24 hrs; do Not freeze. If not used immediately following reconstitution, refrigerate solution and use within 3 hours; do not freeze or shake solution. Store at room temp. Keep in foil pouch until ready to use to protect from light and moisture. | No                                   | Yes                       |               |
| Extavia        | Multiple Sclerosis                      | Injection SubQ            | Yes           | Yes                     | No  | Don't freeze, heat, or shake. Store at 2°C to 25°C (36°F to 77°F). Keep in original carton to protect from light.  | No                                   | Yes                       |               |
| Firmagon       | Hormonal Therapies                      | Injection SubQ            | No            | No                      | Yes   | Don't freeze, heat, or shake. Store at 2°C to 25°C (36°F to 77°F). Keep in original carton to protect from light.  | No                                   | Yes                       |               |
| Flebogamma     | Immune Deficiencies & Related Disorders | IV                        | No            | No                      | Yes   | Don't freeze, heat, or shake. Store at 2°C to 25°C (36°F to 77°F). Keep in original carton to protect from light.  | No                                   | Yes                       |               |
| Flebogamma DIF | Immune Deficiencies & Related Disorders | IV                        | No            | No                      | Yes   | Refrigerate. Avoid excessive or vigorous agitation; do Not shake. Discard if intact vial is left at room temperature for >12 hours prior to use.   | No                                   | Yes                       |               |
| Follistim AQ   | Infertility                             | Injection SubQ            | Yes           | Yes                     | Yes   | Refrigerate; do Not freeze (discard if freezing occurs). Protect from light. Discard pen 28 days after first injection, even if it still contains some unused solution. Do Not use if solution is cloudy, colored, or contains solid particles.  | No                                   | Yes                       |               |
| Forteo         | Osteoporosis                            | Injection SubQ            | Yes           | Yes                     | Yes   | Protect from light<br>Refrigerated at 2°C to 8°C; do Not freeze. The carton of ancillary supplies should be stored at 25°C. After dispensing, store vials at 25°C; once dispensed, vials must be used within 90 days. Once reconstituted, store at <25°C; do Not shake or freeze; use within 3hr   | No                                   | Yes                       |               |
| Fuzeon         | HIV Medications                         | Injection SubQ            | Yes           | Yes                     | No  |  | do not freeze                        | No                        | Yes           |
| Ganirelix      | Infertility                             | Injection SubQ            | Yes           | Yes                     | No  |  | Store below 25°C; do not freeze      | No                        | Yes (generic) |
| Gattex         | Gastrointestinal Disorders- Other       | Injection SubQ            | Yes           | No                      | Yes   | After dispensing, store vials at 25°C; once dispensed, vials must be used within 90 days. Once reconstituted, store at <25°C; do Not shake or freeze; use within 3hr   | No                                   | No                        |               |
| Gel-One        | Osteoarthritis                          | Injection Intra-articular | No            | No                      | Yes   | Store below 25°C; do not freeze  | No                                   | No                        |               |
| Gelsyn-3       | Osteoarthritis                          | Injection Intra-articular | No            | No                      | No  |  | Do not freeze                        | No                        | No            |
| Gemzar         | Oncology- Injectable                    | IV                        | No            | No                      | Yes   | Refrigerate  | No                                   | Yes                       |               |
| GeNotropin     | Growth Hormone & Related Disorders      | Injection SubQ            | Yes           | Yes                     | Yes   | Protect from moisture  | No                                   | No                        |               |
| Gilenya        | Multiple Sclerosis                      | Oral                      | Yes           | No                      | Yes   | Dispense in original bottle; protect from high humidity and light.   | No                                   | Yes                       |               |
| Gilotrif       | Oncology- Oral/Topical                  | Oral                      | Yes           | No                      | Yes   |  | No                                   | No                        |               |

| Drug                          | Treatment For   | Dosage Form               | Covered as Rx | Self-Injectable Product | Costs \$600 or more for up to a 30 day supply | Special Handling (Refrigerate, Limited Distribution, etc)  | Limited Distribution by Manufacturer | Available from Wholesaler |
|-------------------------------|---|---------------------------|---------------|-------------------------|---|--|--------------------------------------|---------------------------|
| Glatiramer acetate (Copaxone) | Multiple Sclerosis  | Injection SubQ            | Yes           | Yes                     | No  | Refrigerate. If needed, may store at 15°C to 30°C for up to 1 month. Avoid exposure to high temperatures; protect from intense light. Do not freeze. Discard if syringe freezes  | No                                   | Yes                       |
| Gonal-F                       | Infertility   | Injection SubQ            | Yes           | Yes                     | Yes   | Store powder refrigerated or at room temperature. Protect from light; do not freeze. Following reconstitution, multidose vials may be stored under refrigeration or at room temperature for up to 28 days.   | No                                   | Yes                       |
| Granix                        | Neutropenia   | Injection SubQ            | Yes           | Yes                     | No  | Refrigerate prefilled syringes. Protect from light. Do not shake. May be removed from for a single period of up to 5 days. If not used within 5 days, the product may be returned to refrigerator up to the expiration date. Dispose of syringes if stored at room temperature for more than 5 days. Exposure to -1°C to -5°C for up to 72 hours and temperatures as low as -15°C to -25°C for up to 24 hours do not adversely affect stability. Discard unused product. | No                                   | Yes                       |
| Harvoni                       | Hepatitis C   | Oral                      | Yes           | No                      | Yes   | Dispense in original container   | No                                   | Yes                       |
| Humantrope                    | Growth Hormone & Related Disorders                            | Injection SubQ            | Yes           | Yes                     | Yes   | Protect from light. Refrigerate  | No                                   | No                        |
| Humira                        | Inflammatory Bowel Disease / Psoriasis / Rheumatoid Arthritis | Injection SubQ            | Yes           | Yes                     | Yes   | Store at 2°C to 8°C (36°F to 46°F) in original container to protect from light; do Not freeze. Do Not use if frozen even if it has been thawed. Do Not store in extreme heat or cold. If needed, may be stored at room temperature up to a maximum of 25°C (77°F) for up to 14 days; discard if Not used within 14 days. Limited Distribution product distributed by CVS Specialty. Store below 30°C; do not freeze. Store in original package to protect from light.    | No                                   | Yes                       |
| Hyalgan                       | Osteoarthritis  | Injection Intra-articular | No            | No                      | No  | Store below 25°C; do not freeze. Store Hymovis in their original packages  | No                                   | Yes                       |
| Hymovis                       | Osteoarthritis  | Injection Intra-articular | No            | No                      | No  |  | No                                   | No                        |
| HyperHEP B                    | Immune Deficiencies & Related Disorders                       | Injection IM              | No            | No                      | No  | Refrigerate. Don't freeze or shake. May be exposed to room temperature for a cumulative 7 days   | No                                   | Yes                       |
| HyppreRHO S/D                 | Immune Deficiencies & Related Disorders                       | Injection IM              | No            | No                      | No  | Store at 2°C to 8°C; do not freeze.  | No                                   | No                        |
| Imantinib (Gleevec)           | Oncology- Oral/Topical  | Oral                      | Yes           | No                      | Yes   | Protect from moisture  | No                                   | Yes                       |

| Drug                        | Treatment For   | Dosage Form        | Covered as Rx | Self-Injectable Product | Costs \$600 or more for up to a 30 day supply | Special Handling (Refrigerate, Limited Distribution, etc)   | Limited Distribution by Manufacturer | Available from Wholesaler |
|-----------------------------|---|--------------------|---------------|-------------------------|---|---|--------------------------------------|---------------------------|
| Increlex (IGF-1 Deficiency) | Growth Hormone & Related Disorders                            | Injection SubQ     | Yes           | Yes                     | Yes   | Protect from light. Refrigerate   | No                                   | No                        |
| Inflextra                   | Inflammatory Bowel Disease / Psoriasis / Rheumatoid Arthritis | IV                 | No            | No                      | Yes   | Refrigerate intact vials at 2°C to 8°C. Manufacturer recommends that solutions diluted in NS for infusion should be used within 3 hours of preparation. However, a stability study of infliximab 0.4 mg/mL prepared in NS in polyvinyl chloride (PVC) bags found no loss of biological activity when stored refrigerated at 4°C for up to 14 days   | No                                   | Yes                       |
| Kevzara                     | Rheumatoid Arthritis  | Injection SubQ     | Yes           | Yes                     | Yes   | Refrigerate, Protect from light. Do Not freeze or shake. After removal from the refrigerator, use within 14 days or discard.  | No                                   | Yes                       |
| Kineret                     | Cryopyrin-Associated Periodic Syndrome / Rheumatoid Arthritis | Injection SubQ     | Yes           | Yes                     | Yes   | Refrigerate. Do Not freeze. Do Not shake. Protect from light. Discard any unused portion.   | No                                   | Yes                       |
| Kisqali                     | Oncology- Oral/Topical  | Oral               | Yes           | No                      | Yes   | Store in the original package   | No                                   | Yes                       |
| Kitabis Pak                 | Cystic Fibrosis   | Nebulized solution | Yes           | No                      | No  | Store under refrigeration at 2°C to 8°C. May be stored in foil pouch (opened or unopened) at room temperature of 25°C for up to 28 days. Protect from light. The colorless to pale yellow solution may darken over time if Not stored under refrigeration; however, the color change does Not affect product quality. Do Not use if solution has been stored at room temperature for >28 days. Diluted solution may be stored up to 4 hours at 2°C to 8°C. Diluted solution is also stable for 4 hours at room temperature of 20°C to 25°C; Refrigeration is preferred. The diluted solution should be protected from light, Not frozen, and used within 4 hours of dilution. | No                                   | No                        |
| Krystexxa                   | Gout  | IV                 | No            | No                      | Yes   | Store at 25°C (77°F); excursions permitted between 15°C to 30°C (59°F to 86°F)  | No                                   | No                        |
| Kyleena                     | Contraceptives  | IUD                | No            | No                      | Yes   |   | No                                   | No                        |

| Drug                        | Treatment For                              | Dosage Form               | Covered as Rx | Self-Injectable Product | Costs \$600 or more for up to a 30 day supply | Special Handling (Refrigerate, Limited Distribution, etc)   | Limited Distribution by Manufacturer | Available from Wholesaler |
|-----------------------------|--|---------------------------|---------------|-------------------------|---|---|--------------------------------------|---------------------------|
| Lartruvo                    | Oncology- Injectable                       | IV                        | No            | No                      | Yes   | Store intact vials at 2°C to 8°C; do not freeze. Keep in original carton to protect from light. Do not shake. Solutions diluted for infusion may be stored for up to 24 hours refrigerated and for an additional 4 hours at room temperature (infusion must be completed within this time frame). If refrigerated, allow infusion solution to reach room temperature prior to administration. | No                                   | No                        |
| Leukine                     | Neutropenia                                | Injection SubQ or IV      | Yes           | Yes                     | No  | Refrigerate. Don't freeze or shake  | No                                   | Yes                       |
| Leuprolide acetate (Lupron) | Hormonal Therapies                         | Injection SubQ or IM      | Yes           | Yes                     | Yes   | Room temp   | No                                   | Yes                       |
| Lonsurf                     | Oncology- Oral/Topical                     | Oral                      | Yes           | No                      | Yes   | If stored outside the original bottle, discard Orals after 30 days  | No                                   | No                        |
| Lupaneta Pack               | Hormonal Therapies                         | Injection IM and Oral     | No            | No                      | Yes   | Room temp   | No                                   | Yes                       |
| Lupron Depot                | Hormonal Therapies                         | Injection IM              | No            | No                      | Yes   | Room temp   | No                                   | Yes                       |
| Lynparza                    | Oncology- Oral/Topical                     | Oral                      | Yes           | No                      | Yes   | Orals: Store at 25°C; excursions permitted to 15°C to 30°C. Do not expose Orals to temperatures >40°C. Orals: Store at 20°C to 25°C; excursions permitted to 15°C to 30°C. Protect from moisture.   | No                                   | No                        |
| MeNopur                     | Infertility                                | Injection SubQ            | Yes           | Yes                     | No  | Refrigerate or room temp. Protect from light  | No                                   | Yes                       |
| Micrhogam                   | Immune Deficiencies & Related Disorders    | Injection IM              | No            | No                      | No  | Store at 2°C to 8°C; do not freeze.   | No                                   | No                        |
| MoNovisc                    | Osteoarthritis                             | Injection Intra-articular | No            | No                      | Yes   | Store below 25°C ]; do not freeze. Protect from light   | No                                   | Yes                       |
| Mozobil                     | Hematopoietics                             | Injection SubQ            | No            | No                      | No  | Limited Distribution product distributed by CVS Specialty   | No                                   | No                        |
| Myobloc                     | Batten Disease                             | Injection IM              | No            | No                      | No  | Refrigerate, Protect from light, once diluted, use within 4 hr  | No                                   | No                        |
| Nabi-HB                     | Immune Deficiencies & Related Disorders    | Injection IM              | No            | No                      | No  | Refrigerate. Do not shake. Use within 6 hours of entering vial.   | No                                   | No                        |
| Neulasta                    | Neutropenia                                | Injection SubQ            | Yes           | Yes                     | Yes   | Refrigerate. Don't freeze or shake. Protect from light  | No                                   | Yes                       |
| Neupogen                    | Neutropenia                                | Injection SubQ or IV      | Yes           | Yes                     | No  | Refrigerate. Store in the original carton. Protect from light. Protect from direct sunlight. Avoid freezing. manufacturer recommends using within 24 hours due to the potential for bacterial contamination.  | No                                   | Yes                       |
| Noritropin                  | Growth Hormone & Related Disorders         | Injection SubQ            | Yes           | Yes                     | Yes   | Refrigerate, but do Not freeze. Protect from light  | No                                   | No                        |
| Nplate                      | Immune (Idiopathic) Thrombocytopenicpurpua | Injection SubQ            | No            | No                      | Yes   | Refrigerate. Protect from light. Store in original carton until use.  | No                                   | Yes                       |
| Nutropin AQ                 | Growth Hormone & Related Disorders         | Injection SubQ            | Yes           | Yes                     | Yes   | Store unopened vials in the refrigerator, away from heat and direct light. Do not freeze. Should be used within 28 days.  | No                                   | No                        |

| Drug                             | Treatment For                      | Dosage Form               | Covered as Rx | Self-Injectable Product | Costs \$600 or more for up to a 30 day supply | Special Handling (Refrigerate, Limited Distribution, etc)  | Limited Distribution by Manufacturer | Available from Wholesaler |
|----------------------------------|------------------------------------|---------------------------|---------------|-------------------------|---|--|--------------------------------------|---------------------------|
| Ocrevus                          | Multiple Sclerosis                 | IV                        | No            | No                      | Yes   | Refrigerate in the outer carton to protect from light. Do not freeze. Do not shake. Solutions diluted for infusion should be used immediately, however may be stored for up to 24 hours refrigerated and 8 hours at room temperature (including infusion time); discard if infusion cannot be completed on the same day. | No                                   | No                        |
| Octreotide acetate (Sandostatin) | Acromegaly                         | Injection SubQ or IV      | Yes           | Yes                     | Yes   | Refrigerate  | No                                   | Yes                       |
| Odomzo                           | Oncology- Oral/Topical             | Oral                      | Yes           | No                      | Yes   | none   | No                                   | No                        |
| Olysio                           | Hepatitis C                        | Oral                      | Yes           | No                      | Yes   | Store in original bottle and protect from light. Protect from light.   | No                                   | Yes                       |
| Omnitrope                        | Growth Hormone & Related Disorders | Injection SubQ            | Yes           | Yes                     | Yes   | Refrigerate  | No                                   | No                        |
| Orenica                          | Rheumatoid Arthritis               | Injection SubQ or IV      | Yes           | Yes                     | Yes   | Refrigerate, protect from light, must used within 24 hr of reconstituting  | No                                   | No                        |
| Orthovisc                        | Osteoarthritis                     | Injection Intra-articular | No            | No                      | No  | Store below 25°C ]; do not freeze. Protect from light  | No                                   | Yes                       |
| Otrexup                          | Psoriasis / Rheumatoid Arthritis   | Injection SubQ            | Yes           | Yes                     | No  | Store intact vials and autoinjectors between 20°C and 25°C (68°F and 77°F); excursions may be permitted between 15°C and 30°C (59°F and 86°F). Protect from light. Refrigerate or room temp for 30 days. Protect from light  | No                                   | Yes                       |
| Ovidrel                          | Infertility                        | Injection SubQ            | Yes           | Yes                     | No  | ed with D5W to a final conc  | No                                   | Yes                       |
| Oxaliplatin                      | Oncology- Injectable               | IV                        | No            | No                      | No  | Refrigerate and protecting from light Intact vial: May be stored at room temperature for up to 14 days   | No                                   | Yes                       |
| Pegasys                          | Hepatitis C                        | Injection SubQ            | Yes           | No                      | Yes   | Prefilled syringe: May be stored at room temperature for up to 6 days  | No                                   | Yes                       |
| Praluent                         | Lipid Disorders- PSK9 Inhibitors   | Injection SubQ            | Yes           | Yes                     | Yes   | Refrigerate in the outer carton to protect from light. Time out of refrigeration should Not exceed 24 hours at 25°C. Do Not freeze. Do Not expose to extreme heat. Do Not shake.   | No                                   | Yes                       |
| Prialt                           | Pain Management                    | Injection Intrathecal     | No            | No                      | Yes   | Prior to use, refrigerate. Once diluted, may be stored at 2°C to 8°C for 24 hours; refrigerate during transit. Do Not freeze. Protect from light.  | No                                   | Yes                       |
| Procrit                          | Anemia                             | Injection SubQ or IV      | Yes           | Yes                     | No  | Vials should be stored at 2°C to 8°C (36°F to 46°F). Do Not freeze. Do Not shake. Protect from light.  | No                                   | Yes                       |



| Drug             | Treatment For                       | Dosage Form        | Covered as Rx | Self-Injectable Product | Costs \$600 or more for up to a 30 day supply | Special Handling (Refrigerate, Limited Distribution, etc)   | Limited Distribution by Manufacturer | Available from Wholesaler |
|------------------|-------------------------------------|--------------------|---------------|-------------------------|---|---|--------------------------------------|---------------------------|
| Prolia           | Osteoporosis<br>Immune (Idiopathic) | Injection SubQ     | No            | No                      | Yes   | Refrigerate in original carton. Do Not freeze. Prior to use, bring to room temperature of 25°C in original container (usually takes 15 to 30 minutes); do Not use any other methods for warming. Use within 14 days once at room temperature. Protect from direct heat and light; do Not expose to temperatures >25°C. Avoid vigorous shaking. Room temp. Dispense in original bottle. Do not remove the infusion bag from the insulated polyurethane container within the shipping box until administration (do not remove the insulated container from the shipping box, or open the lid of the insulated container, until administration). Product may only remain at room temperature for ≤3 hours once removed from shipping container; after removal from shipping container, do not return product to container. | No                                   | Yes                       |
| Promacta         | Thrombocytopenicpurpua              | Oral               | Yes           | No                      | Yes   | Infusion must begin prior to product expiration. Store at 2°C to 8°C in protective foil to protect from light. Refrigerate during transport and do Not expose to room temp for ≥24 hours.   | No                                   | Yes                       |
| Provenge         | Oncology- Injectable                | IV                 | No            | No                      | Yes   | Store intact vials and autoinjectors between 20°C and 25°C (68°F and 77°F); excursions may be permitted between 15°C and 30°C (59°F and 86°F). Protect from light. Refrigerate  | No                                   | No                        |
| Pulmozyme        | Cystic Fibrosis                     | Nebulized Solution | Yes           | No                      | Yes   | Refrigerate; do not freeze. Protect from heat and light. Refrigeration is preferred; however, if needed, may be stored at 2°C to 25°C for up to 30 days.  | No                                   | Yes                       |
| Rasuvo           | Psoriasis / Rheumatoid Arthritis    | Injection SubQ     | Yes           | Yes                     | Yes   |   | No                                   | Yes                       |
| Rebetol Solution | Hepatitis C                         | Oral               | Yes           | No                      | No  |   | No                                   | No                        |
| Rebif            | Multiple Sclerosis                  | Injection SubQ     | Yes           | Yes                     | Yes   |   | No                                   | No                        |

| Drug                               | Treatment For   | Dosage Form          | Covered as Rx | Self-Injectable Product | Costs \$600 or more for up to a 30 day supply | Special Handling (Refrigerate, Limited Distribution, etc)   | Limited Distribution by Manufacturer | Available from Wholesaler |
|------------------------------------|---|----------------------|---------------|-------------------------|---|---|--------------------------------------|---------------------------|
| Remicade                           | Inflammatory Bowel Disease / Psoriasis / Rheumatoid Arthritis | IV                   | No            | No                      | Yes   | Refrigerate intact vials at 2°C to 8°C. Manufacturer recommends that solutions diluted in NS for infusion should be used within 3 hours of preparation. However, a stability study of infliximab 0.4 mg/mL prepared in NS in polyvinyl chloride (PVC) bags found no loss of biological activity when stored refrigerated at 4°C for up to 14 days | No                                   | Yes                       |
| Repatha                            | Lipid Disorders- PSK9 Inhibitors                              | Injection SubQ       | Yes           | Yes                     | Yes   | Refrigerate in the original carton. May also be kept at room temperature in the original carton; however, must use within 30 days (discard if Not used within 30 days). Protect from direct light and do Not expose to temp above 25°C. Do Not freeze. Do Not shake.  | No                                   | Yes                       |
| Repronex                           | Infertility   | Injection SubQ or IM | Yes           | Yes                     | No  | Refrigerate or room temp. Protect from light  | No                                   | Yes                       |
| Rhogam                             | Immune Deficiencies & Related Disorders                       | Injection IM         | No            | No                      | No  | Store at 2°C to 8°C (35°F to 46°F); do not freeze.  | No                                   | Yes                       |
| Rhophylac                          | Immune Deficiencies & Related Disorders                       | Injection IM or IV   | No            | No                      | No  | Store at 2°C to 8°C; do not freeze. Protect from light.   | No                                   | Yes                       |
| Ribapak                            | Hepatitis C   | Oral                 | Yes           | No                      | Yes   | Store at 25°C (77°F); excursions permitted between 15°C and 30°C (59°F and 86°F)  | No                                   | Yes                       |
| Ribasphere                         | Hepatitis C   | Oral                 | Yes           | No                      | No  | None  | No                                   | Yes                       |
| Ribavirin caps (rebetol)           | Hepatitis C   | Oral                 | Yes           | No                      | Yes   | None  | No                                   | Yes                       |
| Ribavirin tabs (Copegus, Moderiba) | Hepatitis C   | Oral                 | Yes           | No                      | Yes   | None  | No                                   | Yes                       |
| Rubraca                            | Oncology- Oral/Topical  | Oral                 | Yes           | No                      | Yes   | none  | No                                   | No                        |
| Rydapt                             | Oncology- Oral/Topical  | Oral                 | Yes           | No                      | Yes   | Store in the original package to protect from moisture.   | No                                   | Yes                       |
| Saizen                             | Growth Hormone & Related Disorders                            | Injection SubQ       | Yes           | Yes                     | Yes   | Protect from light. Refrigerate   | No                                   | Yes                       |
| Samsca                             | Electrolyte Disorders   | Oral                 | Yes           | No                      | Yes   | Store at 25°C; excursions permitted between 15°C and 30°C.  | No                                   | No                        |
| Sandostatin Lar                    | Acromegaly  | Injection IM         | No            | No                      | Yes   | Refrigerate   | No                                   | Yes                       |
| Sensipar                           | Renal Disease   | Oral                 | Yes           | No                      | Yes   | Store at 25°C (77°F); excursions permitted to 15°C to 30°C (59°F to 86°F).  | No                                   | Yes                       |
| Sildenafil (Revatio)               | Pulmonary Arterial Hypertension                               | Oral                 | Yes           | No                      | No  | 86°F).  | No                                   | Yes                       |
| Simponi                            | Inflammatory Bowel Disease / Rheumatoid Arthritis             | Injection SubQ       | Yes           | Yes                     | Yes   | Refrigerate. Do not freeze. Do not shake. Store product in original carton to protect from light.   | No                                   | Yes                       |
| Simponi Aria                       | Rheumatoid Arthritis  | IV                   | No            | No                      | Yes   | Refrigerate; do Not freeze. Do Not shake. Store product in original carton to protect from light  | No                                   | Yes                       |
| Somatuline Depot                   | Acromegaly  | Injection SubQ       | No            | No                      | Yes   | IV: Solutions diluted for infusion may be stored at room temperature for 4 hours.   | No                                   | No                        |

| Drug           | Treatment For                                     | Dosage Form               | Covered as Rx | Self-Injectable Product | Costs \$600 or more for up to a 30 day supply | Special Handling (Refrigerate, Limited Distribution, etc)  | Limited Distribution by Manufacturer | Available from Wholesaler |
|----------------|---|---------------------------|---------------|-------------------------|---|--|--------------------------------------|---------------------------|
| Sovaldi        | Hepatitis C                                       | Oral                      | Yes           | No                      | Yes   | Dispense in original container. Protect from light.  | No                                   | Yes                       |
| Spinraza       | Spinal Muscular atrophy                           | Injection Intrathecal     | No            | No                      | No  | Refrigerate. Discard after 30 hours after being out of original container  | No                                   | No                        |
| Sprycel        | Oncology- Oral/Topical                            | Oral                      | Yes           | No                      | Yes   | none   | No                                   | Yes                       |
| Stelara        | Inflammatory Bowel Disease / Rheumatoid Arthritis | Injection SubQ or IV      | Yes           | Yes                     | Yes   | Refrigerate in original container to protect from light; do not freeze. Do not use if frozen even if it has been thawed. Do not store in extreme heat or cold. If needed, may be stored at room temperature up to a maximum of 25°C for up to 14 days; discard if not used within 14 days.   | No                                   | Yes                       |
| Supartz FX     | Osteoarthritis                                    | Injection Intra-articular | No            | No                      | No  | Store below 30°C (86°C); c   | No                                   | No                        |
| Sutent         | Oncology- Oral/Topical                            | Oral                      | Yes           | No                      | Yes   | none   | No                                   | Yes                       |
| Sylvant        | Oncology- Injectable                              | IV                        | No            | No                      | Yes   | Store intact vials at 2°C to 8°C; protect from light. Reconstituted solution should be further diluted in D5W for infusion within 2 hours; complete infusion within 4 hours of dilution of the reconstituted solution to the infusion container. Discard any unused portion of the reconstituted solution or solution diluted for infusion.      | No                                   | No                        |
| Synagis        | Respiratory Syncytial Virus                       | Injection IM              | No            | No                      | Yes   | Refrigerate in original container. Don't freeze  | No                                   | No                        |
| Synvisc        | Osteoarthritis                                    | Injection Intra-articular | No            | No                      | No  | Store below 30°C (86°C); c   | No                                   | Yes                       |
| Synvisc One    | Osteoarthritis                                    | Injection Intra-articular | No            | No                      | Yes   | Store below 30°C (86°C); c   | No                                   | Yes                       |
| Targretin GelY | Oncology- Oral/Topical                            | Gel                       | Yes           | No                      | Yes   | none   | No                                   | Yes                       |
| Tasigna        | Oncology- Oral/Topical                            | Oral                      | Yes           | No                      | Yes   | none   | No                                   | Yes                       |
| Taxotere       | Oncology- Injectable                              | IV                        | No            | No                      | No  | Store intact vials between 2°C to 25°C. Protect from bright light. Freezing does not adversely affect the product. Solutions diluted for infusion in D5W or NS in non-PVC containers should be used within 6 hours of preparation, including infusion time, when stored between 2°C to 25°C or within 48 hours when stored between 2°C to 8°C    | No                                   | Yes                       |
| Tecentriq      | Oncology- Injectable                              | IV                        | No            | No                      | Yes   | Refrigerate. Do not freeze. Do not shake. Store in original carton to protect from light. Solutions diluted in NS for infusion should be used immediately after preparation; if not used immediately, may be stored for up to 6 hours (including administration time) at room temperature or 24 hours refrigerated at 2°C to 8°C. Do not freeze. | No                                   | No                        |

| Drug                                  | Treatment For                                   | Dosage Form            | Covered as Rx | Self-Injectable Product | Costs \$600 or more for up to a 30 day supply | Special Handling (Refrigerate, Limited Distribution, etc)   | Limited Distribution by Manufacturer | Available from Wholesaler |
|---------------------------------------|---|------------------------|---------------|-------------------------|---|---|--------------------------------------|---------------------------|
| Technivie                             | Hepatitis C                                     | Oral                   | Yes           | No                      | Yes   | Dispense in original container<br>Store intact vials refrigerated at 2°C to 8°C. Reconstituted vials may be stored for up to 14 hours at room temperature of 25°C; infusion must be completed within 14 hours of reconstitution.  | No                                   | Yes                       |
| Temodar<br>Temozolomide (Temodar)     | Oncology- Injectable<br>Oncology- Oral/Topical  | Oral or IV<br>Oral     | No<br>Yes     | No<br>No                | Yes<br>Yes                                    | none<br>Prior to reconstitution, store at 2°C to 8°C (36°F to 46°F). Following reconstitution with the provided diluent, store under refrigeration and use within 14 days (5 mg vial) or 28 days (10 mg vial); do Not freeze. Protect from light. Keep in original package. | No<br>No                             | Yes<br>Yes                |
| Tev-Tropin                            | Growth Hormone & Related Disorders              | Injection SubQ         | Yes           | Yes                     | No  | Protect from light. Keep in original package.   | No                                   | No                        |
| Thalomid                              | Oncology- Oral/Topical                          | Oral                   | Yes           | No                      | Yes   | Store in original package at 25°C (77°F); excursions permitted to 15°C to 30°C (59°F to 86°F). Protect from moisture.   | No                                   | Yes                       |
| Tobramycin inhalation solution (Tobi) | Cystic Fibrosis                                 | Nebulized Solution     | Yes           | No                      | No  | Store at 20°C to 25°C. Do Not freeze MIXJECT system. Administer immediately after reconstitution.   | No                                   | No                        |
| Trelstar                              | Hormonal Therapies                              | Injection IM           | No            | No                      | Yes   | Store refrigerated. Do Not freeze or heat. After first use, store for up to 30 days at 20°C to 25°C   | No                                   | No                        |
| Tymlos<br>Upravi                      | Osteoporosis<br>Pulmonary Arterial Hypertension | Injection SubQ<br>Oral | Yes<br>Yes    | Yes<br>No               | Yes<br>Yes                                    | Refrigerate. Keep implant wrapped in the amber pouch for protection from light; do Not freeze. Do Not open implant vial until just before the time of insertion. The implantation insertion kit should be stored at room temperature (do Not refrigerate insertion kit).    | No<br>No                             | No<br>No                  |
| Vantas                                | Hormonal Therapies                              | Implant                | No            | No                      | Yes   | Dispense in original container  | No                                   | No                        |
| Viekira Pak                           | Hepatitis C                                     | Oral                   | Yes           | No                      | Yes   | Dispense in original container  | No                                   | Yes                       |
| Viekira XR                            | Hepatitis C                                     | Oral                   | Yes           | No                      | Yes   | Injection: Store unopened kit at 2°C to 8°C (36°F to 46°F). Kit may be kept at room temperature of ≤25°C (77°F) for ≤7 days prior to use; do Not freeze. Following reconstitution of the suspension, administer immediately.  | No                                   | Yes                       |
| Vivitrol                              | Alcohol/Opioid Withdrawal                       | Injection IM           | No            | No                      | Yes   | Oral: Store at 20°C to 25°C (68°F to 77°F). Store at 2°C to 8°C ]; do not freeze.   | No                                   | Yes                       |
| Wunrho SDF                            | Immune Deficiencies & Related Disorders         | Injection IM or IV     | No            | No                      | Yes   |   | No                                   | Yes                       |

| Drug                      | Treatment For        | Dosage Form          | Covered as Rx | Self-Injectable Product | Costs \$600 or more for up to a 30 day supply | Special Handling (Refrigerate, Limited Distribution, etc)  | Limited Distribution by Manufacturer | Available from Wholesaler |
|---------------------------|----------------------|----------------------|---------------|-------------------------|---|--|--------------------------------------|---------------------------|
| Xeljanz                   | Rheumatoid Arthritis | Oral                 | Yes           | No                      | Yes   | Store at room temp   | No                                   | Yes                       |
| Xeljanz XR                | Rheumatoid Arthritis | Oral                 | Yes           | No                      | Yes   | Store at room temp   | No                                   | Yes                       |
| Xeomin                    | Batten Disease       | Injection IM         | No            | No                      | Yes   | After reconstitution, store in refrigerator at 2°C to 8°C (36°F to 46°F) and administer within 24 hours. Vial should only be used for one injection session and one patient. Refrigerate. Do not freeze. Prior to use, bring to room temperature of 25°C in original container (usually takes 15 to 30 minutes); do not use any other methods for warming. Use within 14 days once at room temperature. Protect from direct heat and light; do not expose to temperatures >25°C. Avoid vigorous shaking.   | No                                   | No                        |
| Xgeva                     | Oncology- Injectable | Injection SubQ       | No            | No                      | Yes   | Refrigerate in the original carton. Protect from light. Avoid freezing; if frozen, thaw in the refrigerator before administration. Discard if frozen more than once. Do not shake. Transport via a pneumatic tube has not been studied. Solutions diluted for infusion may be stored at room temperature for up to 24 hours (infusion must be completed within 24 hours of preparation). Keep in original blister pack until time of use; protect from moisture. Store at room temp. Keep in foil pouch until ready to use to protect from light and moisture. | No                                   | Yes                       |
| Zarxio                    | Neutropenia          | Injection SubQ or IV | Yes           | Yes                     | No  | Store at room temperature of 25°C; excursions permitted to 15°C to 30°C. After opening, stable for 24 hours at 2°C to 8°C. If refrigerated, allow the refrigerated solution to reach room temperature before administration  | No                                   | No                        |
| Zepatier                  | Hepatitis C          | Oral                 | Yes           | No                      | Yes   | Store at room temp. Keep in foil pouch until ready to use to protect from light and moisture.  | No                                   | Yes                       |
| Zoladex                   | Hormonal Therapies   | Implant              | No            | No                      | Yes   | Store at room temperature of 25°C; excursions permitted to 15°C to 30°C. After opening, stable for 24 hours at 2°C to 8°C. If refrigerated, allow the refrigerated solution to reach room temperature before administration  | No                                   | Yes                       |
| Zoledronic acid (Reclast) | Osteoporosis         | IV                   | No            | No                      | Yes   | Store at room temperature of 25°C; excursions permitted to 15°C to 30°C. After opening, stable for 24 hours at 2°C to 8°C. If refrigerated, allow the refrigerated solution to reach room temperature before administration  | No                                   | Yes                       |

| Drug                     | Treatment For                      | Dosage Form    | Covered as Rx | Self-Injectable Product | Costs \$600 or more for up to a 30 day supply | Special Handling (Refrigerate, Limited Distribution, etc)  | Limited Distribution by Manufacturer | Available from Wholesaler |
|--------------------------|------------------------------------|----------------|---------------|-------------------------|---|--|--------------------------------------|---------------------------|
| Zoledronic acid (Zometa) | Oncology- Injectable               | IV             | No            | No                      | Yes   | Store concentrate vials and ready-to-use bottles at 25°C; excursions permitted to 15°C to 30°C. Diluted solutions for infusion in D5W or NS which are not used immediately after preparation should be refrigerated at 2°C to 8°C. Infusion of solution must be completed within 24 hours of preparation. The ready-to-use bottles are for single use only; if any preparation is necessary (preparing reduced dosage for patients with renal impairment), the prepared, diluted solution may be refrigerated at 2°C to 8°C if not used immediately. Infusion of solution must be completed within 24 hours of preparation. The previously withdrawn volume from the ready-to-use solution should be discarded; do not store or reuse.<br>(see Tev-Tropin) | No                                   | Yes                       |
| Zolinza                  | Oncology- Oral/Topical             | Oral           | Yes           | No                      | Yes   |  | No                                   | Yes                       |
| Zomacton                 | Growth Hormone & Related Disorders | Injection SubQ | Yes           | Yes                     | Yes   |  | No                                   | No                        |
| Zorbitive                | Growth Hormone & Related Disorders | Injection SubQ | Yes           | Yes                     | Yes   | Store intact vials and diluent at 15°C to 30°C (59°F to 86°F). Following reconstitution with the provided diluent (bacteriostatic water for injection containing benzyl alcohol), may store under refrigeration at 2°C to 8°C (36°F to 46°F) for up to 14 days; do Not freeze. If reconstituted with sterile water for injection, solution should be used immediately.   | No                                   | No                        |
| Zytiga                   | Oncology- Oral/Topical             | Oral           | Yes           | No                      | Yes   |  | none                                 | No                        |