

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Adcirca	Pulmonary Arterial Hypertension	Oral	Yes	No	Yes		No	Yes
Afinitor	Oncology- Oral/Topical	Oral	Yes	No	Yes	Protect from light; protect from moisture. If not used immediately following reconstitution, refrigerate solution at 2°C to 8°C and use within 3 hours; do not freeze or shake solution.	No	Yes
Betaseron	Multiple Sclerosis	Injection SubQ	Yes	Yes	No	Store at 2°C to 25°C (36°F to 77°F). Protect from light. Avoid humidity and high temperatures after opening bottle	No	Yes
Bexarotene (Targretin)	Oncology- Oral/Topical	Oral	Yes	No	Yes	Lyophilized powder may be stored in the refrigerator or at room temperature of 3°C to 25°C. Protect from light. Discard unused portion.	No	Yes
Bravelle	Infertility	Injection SubQ or IM	Yes	Yes	No	Discard unused portion.	No	Yes
Capecitabine (Xeloda)	Oncology- Oral/Topical	Oral	Yes	No	Yes	Lyophilized powder may be stored in the refrigerator or at room temperature of 3°C to 25°C. Protect from light. Discard unused portion.	No	Yes
Cetrotide	Infertility	Injection SubQ	Yes	Yes	No	Store at intact vials at 15°C to 30°C. Following reconstitution, solution is stable when refrigerated for 30 days (Novarel) or 60 days (Pregnyl).	No	Yes
Chorionic gonadotropin (Novarel, Pregnyl)	Infertility	Injection IM	Yes	Yes	No	Store intact vials and syringes at 2°C to 8°C; do Not freeze. Do Not separate contents of carton prior to use. Protect from light. Bring to room temperature prior to administration. Store reconstituted solution for up to 24 hours at 2°C to 8°C (do Not freeze); Do Not leave at room temperature for more than 2 hours prior to administration.	No	Yes
Cimzia	Inflammatory Bowel Disease / Rheumatoid Arthritis	Injection SubQ	Yes	Yes	Yes		No	Yes
Daklinza	Hepatitis C	Oral	Yes	No	Yes		No	Yes
Dofetilide (Tikosyn)	Cardiac Disorders	Oral	Yes	No	No	None	No	Yes

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Enbrel	Psoriasis / Rheumatoid Arthritis	Injection SubQ	Yes	Yes	Yes	Refrigerate at 2°C to 8°C. Do Not shake. Do Not freeze or store in extreme heat or cold. Store in the original carton to protect from light or physical damage. Individual autoinjectors, prefilled syringes, dose trays (containing multi-use vials and diluent syringes), or prefilled pens may be stored between 20°C and 25°C (68°F and 77°F) for a maximum single period of 14 days (Enbrel) with protection from light and sources of heat and humidity. Once an autoinjector, prefilled syringe, dose tray, or prefilled pen has been stored at room temperature, it should Not be placed back into the refrigerator; discard after 14 days (Enbrel). Dispense in original container.	No	Yes
Epclusa	Hepatitis C	Oral	Yes	No	Yes		No	Yes
Epogen	Anemia	Injection SubQ or IV	Yes	Yes	Yes	Vials should be stored at 2°C to 8°C (36°F to 46°F). Do Not freeze. Do Not shake. Protect from light. If not used immediately following reconstitution, refrigerate solution and use within 3 hours; do not freeze or shake solution. Refridgerate. Avoid excessive or vigorous agitation; do Not shake. Discard if intact vial is left at room temperature for >12 hours prior to use.	No	Yes
Extavia	Multiple Sclerosis	Injection SubQ	Yes	Yes	No		No	Yes
Follistim AQ	Infertility	Injection SubQ	Yes	Yes	Yes	Refrigerate; do Not freeze (discard if freezing occurs). Protect from light. Discard pen 28 days after first injection, even if it still contains some unused solution. Do Not use if solution is cloudy, colored, or contains solid particles.	No	Yes
Forteo	Osteoporosis	Injection SubQ	Yes	Yes	Yes		No	Yes
Fuzeon	HIV Medications	Injection SubQ	Yes	Yes	No		No	Yes
Ganirelix	Infertility	Injection SubQ	Yes	Yes	No		No	Yes
Gilenya	Multiple Sclerosis	Oral	Yes	No	Yes	Protect from light. Protect from moisture. Refrigerate. If needed, may store at 15°C to 30°C for up to 1 month. Avoid exposure to high temperatures; protect from intense light. Do not freeze. Discard if syringe freezes.	No	Yes
Glatiramer acetate (Copaxone)	Multiple Sclerosis	Injection SubQ	Yes	Yes	No		No	Yes

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Gonal-F	Infertility	Injection SubQ	Yes	Yes	Yes	Store powder refrigerated or at room temperature. Protect from light; do not freeze. Following reconstitution, multidose vials may be stored under refrigeration or at room temperature for up to 28 days. Refrigerate prefilled syringes. Protect from light. Do not shake. May be removed from for a single period of up to 5 days. If not used within 5 days, the product may be returned to refrigerator up to the expiration date. Dispose of syringes if stored at room temperature for more than 5 days. Exposure to -1°C to -5°C for up to 72 hours and temperatures as low as -15°C to -25°C for up to 24 hours do not adversely affect stability. Discard unused product. Dispense in original container	No	Yes
Granix	Neutropenia	Injection SubQ	Yes	Yes	No		No	Yes
Harvoni	Hepatitis C	Oral	Yes	No	Yes		No	Yes
Humira	Inflammatory Bowel Disease / Psoriasis / Rheumatoid Arthritis Oncology- Oral/Topical	Injection SubQ	Yes	Yes	Yes	Store at 2°C to 8°C (36°F to 46°F) in original container to protect from light; do Not freeze. Do Not use if frozen even if it has been thawed. Do Not store in extreme heat or cold. If needed, may be stored at room temperature up to a maximum of 25°C (77°F) for up to 14 days; discard if Not used within 14 days. Protect from moisture. Refrigerate, Protect from light. Do Not freeze or shake. After removal from the refrigerator, use within 14 days or discard.	No	Yes
Imatinib (Gleevec)		Oral	Yes	No	Yes		No	Yes
Kevzara	Rheumatoid Arthritis	Injection SubQ	Yes	Yes	Yes		No	Yes
Kineret	Cryopyrin-Associated Periodic Syndrome / Rheumatoid Arthritis	Injection SubQ	Yes	Yes	Yes	Refrigerate. Do Not freeze. Do Not shake. Protect from light. Discard any unused portion. Store in the original package	No	Yes
Kisqali	Oncology- Oral/Topical	Oral	Yes	No	Yes	Refrigerate. Don't freeze or shake	No	Yes
Leukine	Neutropenia	Injection SubQ or IV	Yes	Yes	No	Room temp	No	Yes
Leuprolide acetate (Lupron)	Hormonal Therapies	Injection SubQ or IM	Yes	Yes	Yes	Refrigerate or room temp. Protect from light	No	Yes
MeNopur	Infertility	Injection SubQ	Yes	Yes	No	Refrigerate. Don't freeze or shake. Protect from light	No	Yes
Neulasta	Neutropenia	Injection SubQ	Yes	Yes	Yes		No	Yes

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Neupogen	Neutropenia	Injection SubQ or IV	Yes	Yes	No	Refrigerate. Store in the original carton. Protect from light. Protect from direct sunlight. Avoid freezing. manufacturer recommends using within 24 hours due to the potential for bacterial contamination.	No	Yes
Octreotide acetate (Sandostatin)	Acromegaly	Injection SubQ or IV	Yes	Yes	Yes	Refrigerate	No	Yes
Olysio	Hepatitis C	Oral	Yes	No	Yes	Store in original bottle and protect from light	No	Yes
Otrexup	Psoriasis / Rheumatoid Arthritis	Injection SubQ	Yes	Yes	No	Store intact vials and autoinjectors between 20°C and 25°C (68°F and 77°F); excursions may be permitted between 15°C and 30°C (59°F and 86°F). Protect from light.	No	Yes
Ovidrel	Infertility	Injection SubQ	Yes	Yes	No	Refrigerate or room temp for 30 days. Protect from light	No	Yes
Praluent	Lipid Disorders- PSK9 Inhibitors	Injection SubQ	Yes	Yes	Yes	Refrigerate in the outer carton to protect from light. Time out of refrigeration should Not exceed 24 hours at 25°C. Do Not freeze. Do Not expose to extreme heat. Do Not shake.	No	Yes
Procrit	Anemia Immune (Idiopathic)	Injection SubQ or IV	Yes	Yes	No	Vials should be stored at 2°C to 8°C (36°F to 46°F). Do Not freeze. Do Not shake. Protect from light. Room temp. Dispense in original bottle.	No	Yes
Promacta	Thrombocytopenicpurpua	Oral	Yes	No	Yes	Store at 2°C to 8°C in protective foil to protect from light. Refrigerate during transport and do Not expose to room temp for ≥24 hours.	No	Yes
Pulmozyme	Cystic Fibrosis	Nebulized Solution	Yes	No	Yes	Store intact vials and autoinjectors between 20°C and 25°C (68°F and 77°F); excursions may be permitted between 15°C and 30°C (59°F and 86°F). Protect from light.	No	Yes
Rasuvo	Psoriasis / Rheumatoid Arthritis	Injection SubQ	Yes	Yes	Yes	Refrigerate in the original carton. May also be kept at room temperature in the original carton; however, must use within 30 days (discard if Not used within 30 days). Protect from direct light and do Not expose to temp above 25°C. Do Not freeze. Do Not shake.	No	Yes
Repatha	Lipid Disorders- PSK9 Inhibitors	Injection SubQ	Yes	Yes	Yes	Refrigerate or room temp. Protect from light	No	Yes
Repronex	Infertility	Injection SubQ or IM	Yes	Yes	No	Store at 25°C (77°F); excursions permitted between 15°C and 30°C (59°F and 86°F)	No	Yes
Ribapak	Hepatitis C	Oral	Yes	No	Yes	None	No	Yes
Ribasphere	Hepatitis C	Oral	Yes	No	No	None	No	Yes

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Ribavirin caps (rebetol)	Hepatitis C	Oral	Yes	No	Yes	None	No	Yes
Ribavirin tabs (Copegus, Moderiba)	Hepatitis C	Oral	Yes	No	Yes	None	No	Yes
Rydapt	Oncology- Oral/Topical	Oral	Yes	No	Yes	Store in the original package to protect from moisture.	No	Yes
Saizen	Growth Hormone & Related Disorders	Injection SubQ	Yes	Yes	Yes	Protect from light. Refrigerate	No	Yes
Sensipar	Renal Disease	Oral	Yes	No	Yes	Store at 25°C (77°F); excursions permitted to 15°C to 30°C (59°F to 86°F).	No	Yes
Sildenafil (Revatio)	Pulmonary Arterial Hypertension	Oral	Yes	No	No	Refrigerate. Do not freeze. Do not shake. Store product in original carton to protect from light.	No	Yes
Simponi	Inflammatory Bowel Disease / Rheumatoid Arthritis	Injection SubQ	Yes	Yes	Yes	Dispense in original container	No	Yes
Sovaldi	Hepatitis C	Oral	Yes	No	Yes	none	No	Yes
Sprycel	Oncology- Oral/Topical	Oral	Yes	No	Yes	Refrigerate in original container to protect from light; do not freeze. Do not use if frozen even if it has been thawed. Do not store in extreme heat or cold. If needed, may be stored at room temperature up to a maximum of 25°C for up to 14 days; discard if not used within 14 days.	No	Yes
Stelara	Inflammatory Bowel Disease / Rheumatoid Arthritis	Injection SubQ or IV	Yes	Yes	Yes	none	No	Yes
Sutent	Oncology- Oral/Topical	Oral	Yes	No	Yes	none	No	Yes
Targretin GelY	Oncology- Oral/Topical	Gel	Yes	No	Yes	none	No	Yes
Tasigna	Oncology- Oral/Topical	Oral	Yes	No	Yes	Dispense in original container	No	Yes
Technivie	Hepatitis C	Oral	Yes	No	Yes	none	No	Yes
Temozolomide (Temodar)	Oncology- Oral/Topical	Oral	Yes	No	Yes	Protect from light. Keep in original package.	No	Yes
Thalomid	Oncology- Oral/Topical	Oral	Yes	No	Yes	Dispense in original container	No	Yes
Viekira Pak	Hepatitis C	Oral	Yes	No	Yes	Dispense in original container	No	Yes
Viekira XR	Hepatitis C	Oral	Yes	No	Yes	Store at room temp	No	Yes
Xeljanz	Rheumatoid Arthritis	Oral	Yes	No	Yes	Store at room temp	No	Yes
Xeljanz XR	Rheumatoid Arthritis	Oral	Yes	No	Yes	Keep in original blister pack until time of use; protect from moisture.	No	Yes
Zepatier	Hepatitis C	Oral	Yes	No	Yes	none	No	Yes
Zolinza	Oncology- Oral/Topical	Oral	Yes	No	Yes	none	No	Yes
Zytiga	Oncology- Oral/Topical	Oral	Yes	No	Yes	none	No	Yes