

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Abraxane	Oncology- Injectable	IV	No	No	Yes	Store at room temp. Protect from bright light. Freezing or refrigerating do not adversely affect the stability of intact vials. Different standards apply after reconstitution	No	Yes
Adagen	Immune Deficiencies & Related Disorders	Injection IM	No	No	Yes	Refrigerate; do Not freeze. Discard unused portions; do Not save for further use. Do Not use if previously frozen	No	No
Adcirca	Pulmonary Arterial Hypertension	Oral	Yes	No	Yes		No	Yes
Afinitor	Oncology- Oral/Topical	Oral	Yes	No	Yes	Protect from light; protect from moisture. Reconstituted solution and infusion solution are stable for 24 hours when refrigerated at 2°C to 8°C	No	Yes
Alimta	Oncology- Injectable	IV	No	No	Yes	none	No	Yes
Alunbrig	Oncology- Oral/Topical	Oral	Yes	No	Yes		No	No
Aranesp	Anemia	Injection SubQ or IV	Yes	No	Yes	Store at 2°C to 8°C (36°F to 46°F); do Not freeze. Do Not shake. Protect from light. Store in original carton until use. The following stability information has also been reported: May be stored at room temperature for up to 7 days	No	Yes
Avonex	Multiple Sclerosis	Injection IM	Yes	Yes	Yes	Refrigerate, Do not freeze. Protect from light. Vial: If refrigeration is not available, may be stored at 25°C (77°F) for up to 30 days	No	No
Azacitidine (Vidaza)	Oncology- Injectable	Injection SubQ or IV	Yes	Yes	Yes	Prior to reconstitution, store intact vials at room temperature of 25°C; excursions permitted to 15°C to 30°C	No	No
Benlysta	Systemic Lupus Erythematosus	Injection SubQ or IV	No	No	Yes	Refrigerate. Protect from light. Dispense in original container if applicable. If not used immediately following reconstitution, refrigerate solution at 2°C to 8°C and use within 3 hours; do not freeze or shake solution.	No	Yes
Betaseron	Multiple Sclerosis	Injection SubQ	Yes	Yes	No	Store at 2°C to 25°C (36°F to 77°F). Protect from light. Avoid humidity and high temperatures after opening bottle	No	Yes
Bexarotene (Targretin)	Oncology- Oral/Topical	Oral	Yes	No	Yes	none	No	Yes
Bosulif	Oncology- Oral/Topical	Oral	Yes	No	Yes		No	No

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Botox	Batten Disease	Injection IM	No	No	No	Store unopened vials under refrigeration at 2°C to 8°C (36°F to 46°F) for up to 36 months or until the expiration date on the vial. After reconstitution, store in refrigerator at 2°C to 8°C (36°F to 46°F) and use within 24 hours (does Not contain a preservative). Botox 100-unit vial: The following stability information has also been reported: May be stored at room temperature for up to 5 day Lyophilized powder may be stored in the refrigerator or at room temperature of 3°C to 25°C. Protect from light. Discard unused portion.	No	Yes
Bravelle Capecitabine (Xeloda)	Infertility Oncology- Oral/Topical	Injection SubQ or IM Oral	Yes Yes	Yes No	No Yes	Prior to reconstitution, store at or below 30°C (86°F). Do not freeze. Following reconstitution in a sterile laminar air flow environment, store under refrigeration. Don't freeze, shake, or heat Lyophilized powder may be stored in the refrigerator or at room temperature of 3°C to 25°C. Protect from light. Discard unused portion.	No No	Yes Yes
Carimune NF	Immune Deficiencies & Related Disorders	IV	No	No	Yes	Store intact vials and syringes at 2°C to 8°C; do Not freeze. Do Not separate contents of carton prior to use. Protect from light. Bring to room temperature prior to administration. Store reconstituted solution for up to 24 hours at 2°C to 8°C (do Not freeze); Do Not leave at room temperature for more than 2 hours prior to administration.	No	Yes
Cetrotide	Infertility	Injection SubQ	Yes	Yes	No	Store at intact vials at 15°C to 30°C. Following reconstitution, solution is stable when refrigerated for 30 days (Novarel) or 60 days (Pregnyl).	No	Yes
Chorionic gonadotropin (Novarel, Pregnyl)	Infertility	Injection IM	Yes	Yes	No	Store intact vials and syringes at 2°C to 8°C; do Not freeze. Do Not separate contents of carton prior to use. Protect from light. Bring to room temperature prior to administration. Store reconstituted solution for up to 24 hours at 2°C to 8°C (do Not freeze); Do Not leave at room temperature for more than 2 hours prior to administration.	No	Yes
Cimzia	Inflammatory Bowel Disease / Rheumatoid Arthritis	Injection SubQ	Yes	Yes	Yes	Store at 2°C to 8°C (36°F to 46°F) for up to 36 months or at room temperature for up to 12 months; do Not return vial to refrigerator after it has been stored at room temperature. Do Not freeze. Keep in original carton to protect from light.	No	Yes
Cuvitru	Transplant	Injection SubQ or IM	No	No	Yes	Store at 2°C to 8°C (36°F to 46°F) for up to 36 months or at room temperature for up to 12 months; do Not return vial to refrigerator after it has been stored at room temperature. Do Not freeze. Keep in original carton to protect from light.	No	No
Daklinza	Hepatitis C	Oral	Yes	No	Yes	None	No	Yes
Dofetilide (Tikosyn)	Cardiac Disorders	Oral	Yes	No	No	None	No	Yes

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Dupixent	Atopic Dermatitis	Injection SubQ	Yes	Yes	Yes	N/A	No	No
Dysport	Batten Disease	Injection IM	No	No	Yes	Refrigerate, protect from light	No	No
Eligard (Canadian Brand name, See Lupron)	Hormonal Therapies	Injection SubQ	No	No	Yes	Refrigerate at 2°C to 8°C. Do Not shake. Do Not freeze or store in extreme heat or cold. Store in the original carton to protect from light or physical damage.	No	No
Enbrel	Psoriasis / Rheumatoid Arthritis	Injection SubQ	Yes	Yes	Yes	Individual autoinjectors, prefilled syringes, dose trays (containing multi-use vials and diluent syringes), or prefilled pens may be stored between 20°C and 25°C (68°F and 77°F) for a maximum single period of 14 days (Enbrel) with protection from light and sources of heat and humidity. Once an autoinjector, prefilled syringe, dose tray, or prefilled pen has been stored at room temperature, it should Not be placed back into the refrigerator; discard after 14 days (Enbrel)	No	Yes
Entyvio	Inflammatory Bowel Disease	IV	No	No	Yes	Refrigerate unopened vials. Retain in original package to protect from light. After reconstitution and dilution, the infusion solution may be refrigerated for up to 4 hours). Don't freeze. Discard any unused portion	No	No
Epclusa	Hepatitis C	Oral	Yes	No	Yes	Dispense in original container	No	Yes
Epogen	Anemia	Injection SubQ or IV	Yes	Yes	Yes	Vials should be stored at 2°C to 8°C (36°F to 46°F). Do Not freeze. Do Not shake. Protect from light.	No	Yes
Erwinaze	Oncology- Injectable	Injection IM or IV	No	No	Yes	Refrigerate. Protect from light. Within 15 minutes of reconstitution, withdraw appropriate volume for dose into a polypropylene syringe. Do not freeze or refrigerate reconstituted solution; discard if not administered within 4 hours.	No	No
Euflexxa	Osteoarthritis	Injection Intra-articular	No	No	No	Store refrigerated or at room temperature; do not freeze. Protect from light. If refrigerated, remove from refrigeration at least 20 to 30 minutes before use.	No	Yes

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Exondys	Duchenne Muscular Dystrophy	IV	No	No	Yes	Store at 2°C to 8°C. Do Not freeze. Protect from light and store in the original carton until ready for use. The diluted solution may be stored at 2°C to 8°C for up to 24 hrs; do Not freeze. If not used immediately following reconstitution, refrigerate solution and use within 3 hours; do not freeze or shake solution. Store at room temp. Keep in foil pouch until ready to use to protect from light and moisture.	No	Yes
Extavia	Multiple Sclerosis	Injection SubQ	Yes	Yes	No	Don't freeze, heat, or shake. Store at 2°C to 25°C (36°F to 77°F). Keep in original carton to protect from light.	No	Yes
Firmagon	Hormonal Therapies	Injection SubQ	No	No	Yes	Don't freeze, heat, or shake. Store at 2°C to 25°C (36°F to 77°F). Keep in original carton to protect from light.	No	Yes
Flebogamma	Immune Deficiencies & Related Disorders	IV	No	No	Yes	Don't freeze, heat, or shake. Store at 2°C to 25°C (36°F to 77°F). Keep in original carton to protect from light.	No	Yes
Flebogamma DIF	Immune Deficiencies & Related Disorders	IV	No	No	Yes	Refrigerate. Avoid excessive or vigorous agitation; do Not shake. Discard if intact vial is left at room temperature for >12 hours prior to use.	No	Yes
Follistim AQ	Infertility	Injection SubQ	Yes	Yes	Yes	Refrigerate; do Not freeze (discard if freezing occurs). Protect from light. Discard pen 28 days after first injection, even if it still contains some unused solution. Do Not use if solution is cloudy, colored, or contains solid particles.	No	Yes
Forteo	Osteoporosis	Injection SubQ	Yes	Yes	Yes	Protect from light Refrigerated at 2°C to 8°C; do Not freeze. The carton of ancillary supplies should be stored at 25°C. After dispensing, store vials at 25°C; once dispensed, vials must be used within 90 days. Once reconstituted, store at <25°C; do Not shake or freeze; use within 3hr	No	Yes
Fuzeon	HIV Medications	Injection SubQ	Yes	Yes	No		No	Yes
Ganirelix	Infertility	Injection SubQ	Yes	Yes	No		No	Yes (generic)
Gattex	Gastrointestinal Disorders- Other	Injection SubQ	Yes	No	Yes	Store below 25°C (77°F); do not freeze	No	No
Gel-One	Osteoarthritis	Injection Intra-articular	No	No	Yes	Store below 25°C; do not freeze	No	No
Gelsyn-3	Osteoarthritis	Injection Intra-articular	No	No	No		No	No
Gemzar	Oncology- Injectable	IV	No	No	Yes	Don't freeze	No	Yes
GeNotropin	Growth Hormone & Related Disorders	Injection SubQ	Yes	Yes	Yes	Refrigerate	No	No
Gilenya	Multiple Sclerosis	Oral	Yes	No	Yes	Protect from moisture Dispense in original bottle; protect from high humidity and light.	No	Yes
Gilotrif	Oncology- Oral/Topical	Oral	Yes	No	Yes		No	No

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Glatiramer acetate (Copaxone)	Multiple Sclerosis	Injection SubQ	Yes	Yes	No	Refrigerate. If needed, may store at 15°C to 30°C for up to 1 month. Avoid exposure to high temperatures; protect from intense light. Do not freeze. Discard if syringe freezes	No	Yes
Gonal-F	Infertility	Injection SubQ	Yes	Yes	Yes	Store powder refrigerated or at room temperature. Protect from light; do not freeze. Following reconstitution, multidose vials may be stored under refrigeration or at room temperature for up to 28 days.	No	Yes
Granix	Neutropenia	Injection SubQ	Yes	Yes	No	Refrigerate prefilled syringes. Protect from light. Do not shake. May be removed from for a single period of up to 5 days. If not used within 5 days, the product may be returned to refrigerator up to the expiration date. Dispose of syringes if stored at room temperature for more than 5 days. Exposure to -1°C to -5°C for up to 72 hours and temperatures as low as -15°C to -25°C for up to 24 hours do not adversely affect stability. Discard unused product.	No	Yes
Harvoni	Hepatitis C	Oral	Yes	No	Yes	Dispense in original container	No	Yes
Humantrope	Growth Hormone & Related Disorders	Injection SubQ	Yes	Yes	Yes	Protect from light. Refrigerate	No	No
Humira	Inflammatory Bowel Disease / Psoriasis / Rheumatoid Arthritis	Injection SubQ	Yes	Yes	Yes	Store at 2°C to 8°C (36°F to 46°F) in original container to protect from light; do Not freeze. Do Not use if frozen even if it has been thawed. Do Not store in extreme heat or cold. If needed, may be stored at room temperature up to a maximum of 25°C (77°F) for up to 14 days; discard if Not used within 14 days.	No	Yes
Hyalgan	Osteoarthritis	Injection Intra-articular	No	No	No	Limited Distribution product distributed by CVS Specialty. Store below 30°C; do not freeze. Store in original package to protect from light.	No	Yes
Hymovis	Osteoarthritis	Injection Intra-articular	No	No	No	Store below 25°C; do not freeze. Store Hymovis in their original packages	No	No
HyperHEP B	Immune Deficiencies & Related Disorders	Injection IM	No	No	No	Refrigerate. Don't freeze or shake. May be exposed to room temperature for a cumulative 7 days	No	Yes
HyppreRHO S/D	Immune Deficiencies & Related Disorders	Injection IM	No	No	No	Store at 2°C to 8°C; do not freeze.	No	No
Imantinib (Gleevec)	Oncology- Oral/Topical	Oral	Yes	No	Yes	Protect from moisture	No	Yes

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Increlex (IGF-1 Deficiency)	Growth Hormone & Related Disorders	Injection SubQ	Yes	Yes	Yes	Protect from light. Refrigerate	No	No
Inflextra	Inflammatory Bowel Disease / Psoriasis / Rheumatoid Arthritis	IV	No	No	Yes	Refrigerate intact vials at 2°C to 8°C. Manufacturer recommends that solutions diluted in NS for infusion should be used within 3 hours of preparation. However, a stability study of infliximab 0.4 mg/mL prepared in NS in polyvinyl chloride (PVC) bags found no loss of biological activity when stored refrigerated at 4°C for up to 14 days	No	Yes
Kevzara	Rheumatoid Arthritis	Injection SubQ	Yes	Yes	Yes	Refrigerate, Protect from light. Do Not freeze or shake. After removal from the refrigerator, use within 14 days or discard.	No	Yes
Kineret	Cryopyrin-Associated Periodic Syndrome / Rheumatoid Arthritis	Injection SubQ	Yes	Yes	Yes	Refrigerate. Do Not freeze. Do Not shake. Protect from light. Discard any unused portion.	No	Yes
Kisqali	Oncology- Oral/Topical	Oral	Yes	No	Yes	Store in the original package	No	Yes
Kitabis Pak	Cystic Fibrosis	Nebulized solution	Yes	No	No	Store under refrigeration at 2°C to 8°C. May be stored in foil pouch (opened or uNopened) at room temperature of 25°C for up to 28 days. Protect from light. The colorless to pale yellow solution may darken over time if Not stored under refrigeration; however, the color change does Not affect product quality. Do Not use if solution has been stored at room temperature for >28 days. Diluted solution may be stored up to 4 hours at 2°C to 8°C. Diluted solution is also stable for 4 hours at room temperature of 20°C to 25°C; Refrigeration is preferred. The diluted solution should be protected from light, Not frozen, and used within 4 hours of dilution.	No	No
Krystexxa	Gout	IV	No	No	Yes	Store at 25°C (77°F); excursions permitted between 15°C to 30°C (59°F to 86°F)	No	No
Kyleena	Contraceptives	IUD	No	No	Yes		No	No

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Lartruvo	Oncology- Injectable	IV	No	No	Yes	Store intact vials at 2°C to 8°C; do not freeze. Keep in original carton to protect from light. Do not shake. Solutions diluted for infusion may be stored for up to 24 hours refrigerated and for an additional 4 hours at room temperature (infusion must be completed within this time frame). If refrigerated, allow infusion solution to reach room temperature prior to administration.	No	No
Leukine	Neutropenia	Injection SubQ or IV	Yes	Yes	No	Refrigerate. Don't freeze or shake	No	Yes
Leuprolide acetate (Lupron)	Hormonal Therapies	Injection SubQ or IM	Yes	Yes	Yes	Room temp	No	Yes
Lonsurf	Oncology- Oral/Topical	Oral	Yes	No	Yes	If stored outside the original bottle, discard Orals after 30 days	No	No
Lupaneta Pack	Hormonal Therapies	Injection IM and Oral	No	No	Yes	Room temp	No	Yes
Lupron Depot	Hormonal Therapies	Injection IM	No	No	Yes	Room temp	No	Yes
Lynparza	Oncology- Oral/Topical	Oral	Yes	No	Yes	Orals: Store at 25°C; excursions permitted to 15°C to 30°C. Do not expose Orals to temperatures >40°C. Orals: Store at 20°C to 25°C; excursions permitted to 15°C to 30°C. Protect from moisture.	No	No
MeNopur	Infertility	Injection SubQ	Yes	Yes	No	Refrigerate or room temp. Protect from light	No	Yes
Micrhogam	Immune Deficiencies & Related Disorders	Injection IM	No	No	No	Store at 2°C to 8°C; do not freeze.	No	No
MoNovisc	Osteoarthritis	Injection Intra-articular	No	No	Yes	Store below 25°C]; do not freeze. Protect from light	No	Yes
Mozobil	Hematopoietics	Injection SubQ	No	No	No	Limited Distribution product distributed by CVS Specialty	No	No
Myobloc	Batten Disease	Injection IM	No	No	No	Refrigerate, Protect from light, once diluted, use within 4 hr	No	No
Nabi-HB	Immune Deficiencies & Related Disorders	Injection IM	No	No	No	Refrigerate. Do not shake. Use within 6 hours of entering vial.	No	No
Neulasta	Neutropenia	Injection SubQ	Yes	Yes	Yes	Refrigerate. Don't freeze or shake. Protect from light	No	Yes
Neupogen	Neutropenia	Injection SubQ or IV	Yes	Yes	No	Refrigerate. Store in the original carton. Protect from light. Protect from direct sunlight. Avoid freezing. manufacturer recommends using within 24 hours due to the potential for bacterial contamination.	No	Yes
Noritropin	Growth Hormone & Related Disorders	Injection SubQ	Yes	Yes	Yes	Refrigerate, but do Not freeze. Protect from light	No	No
Nplate	Immune (Idiopathic) Thrombocytopenicpurpua	Injection SubQ	No	No	Yes	Refrigerate. Protect from light. Store in original carton until use.	No	Yes
Nutropin AQ	Growth Hormone & Related Disorders	Injection SubQ	Yes	Yes	Yes	Store unopened vials in the refrigerator, away from heat and direct light. Do not freeze. Should be used within 28 days.	No	No

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Ocrevus	Multiple Sclerosis	IV	No	No	Yes	Refrigerate in the outer carton to protect from light. Do not freeze. Do not shake. Solutions diluted for infusion should be used immediately, however may be stored for up to 24 hours refrigerated and 8 hours at room temperature (including infusion time); discard if infusion cannot be completed on the same day.	No	No
Octreotide acetate (Sandostatin)	Acromegaly	Injection SubQ or IV	Yes	Yes	Yes	Refrigerate	No	Yes
Odomzo	Oncology- Oral/Topical	Oral	Yes	No	Yes	none	No	No
Olysio	Hepatitis C	Oral	Yes	No	Yes	Store in original bottle and protect from light. Protect from light.	No	Yes
Omnitrope	Growth Hormone & Related Disorders	Injection SubQ	Yes	Yes	Yes	Refrigerate	No	No
Orenica	Rheumatoid Arthritis	Injection SubQ or IV	Yes	Yes	Yes	Refrigerate, protect from light, must used within 24 hr of reconstituting	No	No
Orthovisc	Osteoarthritis	Injection Intra-articular	No	No	No	Store below 25°C]; do not freeze. Protect from light	No	Yes
Otrexup	Psoriasis / Rheumatoid Arthritis	Injection SubQ	Yes	Yes	No	Store intact vials and autoinjectors between 20°C and 25°C (68°F and 77°F); excursions may be permitted between 15°C and 30°C (59°F and 86°F). Protect from light. Refrigerate or room temp for 30 days. Protect from light	No	Yes
Ovidrel	Infertility	Injection SubQ	Yes	Yes	No	ed with D5W to a final conc	No	Yes
Oxaliplatin	Oncology- Injectable	IV	No	No	No	Refrigerate and protecting from light Intact vial: May be stored at room temperature for up to 14 days	No	Yes
Pegasys	Hepatitis C	Injection SubQ	Yes	No	Yes	Prefilled syringe: May be stored at room temperature for up to 6 days	No	Yes
Praluent	Lipid Disorders- PSK9 Inhibitors	Injection SubQ	Yes	Yes	Yes	Refrigerate in the outer carton to protect from light. Time out of refrigeration should Not exceed 24 hours at 25°C. Do Not freeze. Do Not expose to extreme heat. Do Not shake.	No	Yes
Prialt	Pain Management	Injection Intrathecal	No	No	Yes	Prior to use, refrigerate. Once diluted, may be stored at 2°C to 8°C for 24 hours; refrigerate during transit. Do Not freeze. Protect from light.	No	Yes
Procrit	Anemia	Injection SubQ or IV	Yes	Yes	No	Vials should be stored at 2°C to 8°C (36°F to 46°F). Do Not freeze. Do Not shake. Protect from light.	No	Yes

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Prolia	Osteoporosis Immune (Idiopathic)	Injection SubQ	No	No	Yes	Refrigerate in original carton. Do Not freeze. Prior to use, bring to room temperature of 25°C in original container (usually takes 15 to 30 minutes); do Not use any other methods for warming. Use within 14 days once at room temperature. Protect from direct heat and light; do Not expose to temperatures >25°C. Avoid vigorous shaking. Room temp. Dispense in original bottle. Do not remove the infusion bag from the insulated polyurethane container within the shipping box until administration (do not remove the insulated container from the shipping box, or open the lid of the insulated container, until administration). Product may only remain at room temperature for ≤3 hours once removed from shipping container; after removal from shipping container, do not return product to container. Infusion must begin prior to product expiration. Store at 2°C to 8°C in protective foil to protect from light. Refrigerate during transport and do Not expose to room temp for ≥24 hours.	No	Yes
Promacta	Thrombocytopenicpurpua	Oral	Yes	No	Yes		No	Yes
Provenge	Oncology- Injectable	IV	No	No	Yes		No	No
Pulmozyme	Cystic Fibrosis	Nebulized Solution	Yes	No	Yes		No	Yes
Rasuvo	Psoriasis / Rheumatoid Arthritis	Injection SubQ	Yes	Yes	Yes	Store intact vials and autoinjectors between 20°C and 25°C (68°F and 77°F); excursions may be permitted between 15°C and 30°C (59°F and 86°F). Protect from light.	No	Yes
Rebetol Solution	Hepatitis C	Oral	Yes	No	No	Refrigerate. Refrigerate; do not freeze. Protect from heat and light. Refrigeration is preferred; however, if needed, may be stored at 2°C to 25°C for up to 30 days.	No	No
Rebif	Multiple Sclerosis	Injection SubQ	Yes	Yes	Yes		No	No

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Remicade	Inflammatory Bowel Disease / Psoriasis / Rheumatoid Arthritis	IV	No	No	Yes	Refrigerate intact vials at 2°C to 8°C. Manufacturer recommends that solutions diluted in NS for infusion should be used within 3 hours of preparation. However, a stability study of infliximab 0.4 mg/mL prepared in NS in polyvinyl chloride (PVC) bags found no loss of biological activity when stored refrigerated at 4°C for up to 14 days	No	Yes
Repatha	Lipid Disorders- PSK9 Inhibitors	Injection SubQ	Yes	Yes	Yes	Refrigerate in the original carton. May also be kept at room temperature in the original carton; however, must use within 30 days (discard if Not used within 30 days). Protect from direct light and do Not expose to temp above 25°C. Do Not freeze. Do Not shake.	No	Yes
Repronex	Infertility	Injection SubQ or IM	Yes	Yes	No	Refrigerate or room temp. Protect from light	No	Yes
Rhogam	Immune Deficiencies & Related Disorders	Injection IM	No	No	No	Store at 2°C to 8°C (35°F to 46°F); do not freeze.	No	Yes
Rhophylac	Immune Deficiencies & Related Disorders	Injection IM or IV	No	No	No	Store at 2°C to 8°C; do not freeze. Protect from light.	No	Yes
Ribapak	Hepatitis C	Oral	Yes	No	Yes	Store at 25°C (77°F); excursions permitted between 15°C and 30°C (59°F and 86°F)	No	Yes
Ribasphere	Hepatitis C	Oral	Yes	No	No	None	No	Yes
Ribavirin caps (rebetol)	Hepatitis C	Oral	Yes	No	Yes	None	No	Yes
Ribavirin tabs (Copegus, Moderiba)	Hepatitis C	Oral	Yes	No	Yes	None	No	Yes
Rubraca	Oncology- Oral/Topical	Oral	Yes	No	Yes	none	No	No
Rydapt	Oncology- Oral/Topical	Oral	Yes	No	Yes	Store in the original package to protect from moisture.	No	Yes
Saizen	Growth Hormone & Related Disorders	Injection SubQ	Yes	Yes	Yes	Protect from light. Refrigerate	No	Yes
Samsca	Electrolyte Disorders	Oral	Yes	No	Yes	Store at 25°C; excursions permitted between 15°C and 30°C.	No	No
Sandostatin Lar	Acromegaly	Injection IM	No	No	Yes	Refrigerate	No	Yes
Sensipar	Renal Disease	Oral	Yes	No	Yes	Store at 25°C (77°F); excursions permitted to 15°C to 30°C (59°F to 86°F).	No	Yes
Sildenafil (Revatio)	Pulmonary Arterial Hypertension	Oral	Yes	No	No	86°F).	No	Yes
Simponi	Inflammatory Bowel Disease / Rheumatoid Arthritis	Injection SubQ	Yes	Yes	Yes	Refrigerate. Do not freeze. Do not shake. Store product in original carton to protect from light.	No	Yes
Simponi Aria	Rheumatoid Arthritis	IV	No	No	Yes	Refrigerate; do Not freeze. Do Not shake. Store product in original carton to protect from light	No	Yes
Somatuline Depot	Acromegaly	Injection SubQ	No	No	Yes	IV: Solutions diluted for infusion may be stored at room temperature for 4 hours.	No	No

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Sovaldi	Hepatitis C	Oral	Yes	No	Yes	Dispense in original container. Protect from light.	No	Yes
Spinraza	Spinal Muscular atrophy	Injection Intrathecal	No	No	No	Refrigerate. Discard after 30 hours after being out of original container	No	No
Sprycel	Oncology- Oral/Topical	Oral	Yes	No	Yes	none	No	Yes
Stelara	Inflammatory Bowel Disease / Rheumatoid Arthritis	Injection SubQ or IV	Yes	Yes	Yes	Refrigerate in original container to protect from light; do not freeze. Do not use if frozen even if it has been thawed. Do not store in extreme heat or cold. If needed, may be stored at room temperature up to a maximum of 25°C for up to 14 days; discard if not used within 14 days.	No	Yes
Supartz FX	Osteoarthritis	Injection Intra-articular	No	No	No	Store below 30°C (86°C); c	No	No
Sutent	Oncology- Oral/Topical	Oral	Yes	No	Yes	none	No	Yes
Sylvant	Oncology- Injectable	IV	No	No	Yes	Store intact vials at 2°C to 8°C; protect from light. Reconstituted solution should be further diluted in D5W for infusion within 2 hours; complete infusion within 4 hours of dilution of the reconstituted solution to the infusion container. Discard any unused portion of the reconstituted solution or solution diluted for infusion.	No	No
Synagis	Respiratory Syncytial Virus	Injection IM	No	No	Yes	Refrigerate in original container. Don't freeze	No	No
Synvisc	Osteoarthritis	Injection Intra-articular	No	No	No	Store below 30°C (86°C); c	No	Yes
Synvisc One	Osteoarthritis	Injection Intra-articular	No	No	Yes	Store below 30°C (86°C); c	No	Yes
Targretin GelY	Oncology- Oral/Topical	Gel	Yes	No	Yes	none	No	Yes
Tasigna	Oncology- Oral/Topical	Oral	Yes	No	Yes	none	No	Yes
Taxotere	Oncology- Injectable	IV	No	No	No	Store intact vials between 2°C to 25°C. Protect from bright light. Freezing does not adversely affect the product. Solutions diluted for infusion in D5W or NS in non-PVC containers should be used within 6 hours of preparation, including infusion time, when stored between 2°C to 25°C or within 48 hours when stored between 2°C to 8°C	No	Yes
Tecentriq	Oncology- Injectable	IV	No	No	Yes	Refrigerate. Do not freeze. Do not shake. Store in original carton to protect from light. Solutions diluted in NS for infusion should be used immediately after preparation; if not used immediately, may be stored for up to 6 hours (including administration time) at room temperature or 24 hours refrigerated at 2°C to 8°C. Do not freeze.	No	No

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Technivie	Hepatitis C	Oral	Yes	No	Yes	Dispense in original container Store intact vials refrigerated at 2°C to 8°C. Reconstituted vials may be stored for up to 14 hours at room temperature of 25°C; infusion must be completed within 14 hours of reconstitution.	No	Yes
Temodar Temozolomide (Temodar)	Oncology- Injectable Oncology- Oral/Topical	Oral or IV Oral	No Yes	No No	Yes Yes	none Prior to reconstitution, store at 2°C to 8°C (36°F to 46°F). Following reconstitution with the provided diluent, store under refrigeration and use within 14 days (5 mg vial) or 28 days (10 mg vial); do Not freeze. Protect from light. Keep in original package.	No No	Yes Yes
Tev-Tropin	Growth Hormone & Related Disorders	Injection SubQ	Yes	Yes	No	Protect from light. Keep in original package.	No	No
Thalomid	Oncology- Oral/Topical	Oral	Yes	No	Yes	Store in original package at 25°C (77°F); excursions permitted to 15°C to 30°C (59°F to 86°F). Protect from moisture.	No	Yes
Tobramycin inhalation solution (Tobi)	Cystic Fibrosis	Nebulized Solution	Yes	No	No	Store at 20°C to 25°C. Do Not freeze MIXJECT system. Administer immediately after reconstitution.	No	No
Trelstar	Hormonal Therapies	Injection IM	No	No	Yes	Store refrigerated. Do Not freeze or heat. After first use, store for up to 30 days at 20°C to 25°C	No	No
Tymlos Uptravi	Osteoporosis Pulmonary Arterial Hypertension	Injection SubQ Oral	Yes Yes	Yes No	Yes Yes	Refrigerate. Keep implant wrapped in the amber pouch for protection from light; do Not freeze. Do Not open implant vial until just before the time of insertion. The implantation insertion kit should be stored at room temperature (do Not refrigerate insertion kit).	No No	No No
Vantas	Hormonal Therapies	Implant	No	No	Yes	Dispense in original container	No	No
Viekira Pak	Hepatitis C	Oral	Yes	No	Yes	Dispense in original container	No	Yes
Viekira XR	Hepatitis C	Oral	Yes	No	Yes	Injection: Store unopened kit at 2°C to 8°C (36°F to 46°F). Kit may be kept at room temperature of ≤25°C (77°F) for ≤7 days prior to use; do Not freeze. Following reconstitution of the suspension, administer immediately.	No	Yes
Vivitrol	Alcohol/Opioid Withdrawal	Injection IM	No	No	Yes	Oral: Store at 20°C to 25°C (68°F to 77°F). Store at 2°C to 8°C]; do not freeze.	No	Yes
Wunrho SDF	Immune Deficiencies & Related Disorders	Injection IM or IV	No	No	Yes		No	Yes

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Xeljanz	Rheumatoid Arthritis	Oral	Yes	No	Yes	Store at room temp	No	Yes
Xeljanz XR	Rheumatoid Arthritis	Oral	Yes	No	Yes	Store at room temp	No	Yes
Xeomin	Batten Disease	Injection IM	No	No	Yes	After reconstitution, store in refrigerator at 2°C to 8°C (36°F to 46°F) and administer within 24 hours. Vial should only be used for one injection session and one patient. Refrigerate. Do not freeze. Prior to use, bring to room temperature of 25°C in original container (usually takes 15 to 30 minutes); do not use any other methods for warming. Use within 14 days once at room temperature. Protect from direct heat and light; do not expose to temperatures >25°C. Avoid vigorous shaking.	No	No
Xgeva	Oncology- Injectable	Injection SubQ	No	No	Yes	Refrigerate in the original carton. Protect from light. Avoid freezing; if frozen, thaw in the refrigerator before administration. Discard if frozen more than once. Do not shake. Transport via a pneumatic tube has not been studied. Solutions diluted for infusion may be stored at room temperature for up to 24 hours (infusion must be completed within 24 hours of preparation). Keep in original blister pack until time of use; protect from moisture. Store at room temp. Keep in foil pouch until ready to use to protect from light and moisture.	No	Yes
Zarxio	Neutropenia	Injection SubQ or IV	Yes	Yes	No	Store at room temperature of 25°C; excursions permitted to 15°C to 30°C. After opening, stable for 24 hours at 2°C to 8°C. If refrigerated, allow the refrigerated solution to reach room temperature before administration	No	No
Zepatier	Hepatitis C	Oral	Yes	No	Yes	Store at room temp. Keep in foil pouch until ready to use to protect from light and moisture.	No	Yes
Zoladex	Hormonal Therapies	Implant	No	No	Yes	Store at room temperature of 25°C; excursions permitted to 15°C to 30°C. After opening, stable for 24 hours at 2°C to 8°C. If refrigerated, allow the refrigerated solution to reach room temperature before administration	No	Yes
Zoledronic acid (Reclast)	Osteoporosis	IV	No	No	Yes	Store at room temperature of 25°C; excursions permitted to 15°C to 30°C. After opening, stable for 24 hours at 2°C to 8°C. If refrigerated, allow the refrigerated solution to reach room temperature before administration	No	Yes

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Zoledronic acid (Zometa)	Oncology- Injectable	IV	No	No	Yes	Store concentrate vials and ready-to-use bottles at 25°C; excursions permitted to 15°C to 30°C. Diluted solutions for infusion in D5W or NS which are not used immediately after preparation should be refrigerated at 2°C to 8°C. Infusion of solution must be completed within 24 hours of preparation. The ready-to-use bottles are for single use only; if any preparation is necessary (preparing reduced dosage for patients with renal impairment), the prepared, diluted solution may be refrigerated at 2°C to 8°C if not used immediately. Infusion of solution must be completed within 24 hours of preparation. The previously withdrawn volume from the ready-to-use solution should be discarded; do not store or reuse. none (see Tev-Tropin)	No	Yes
Zolinza	Oncology- Oral/Topical	Oral	Yes	No	Yes		No	Yes
Zomacton	Growth Hormone & Related Disorders	Injection SubQ	Yes	Yes	Yes		No	No
Zorbitive	Growth Hormone & Related Disorders	Injection SubQ	Yes	Yes	Yes	Store intact vials and diluent at 15°C to 30°C (59°F to 86°F). Following reconstitution with the provided diluent (bacteriostatic water for injection containing benzyl alcohol), may store under refrigeration at 2°C to 8°C (36°F to 46°F) for up to 14 days; do Not freeze. If reconstituted with sterile water for injection, solution should be used immediately. none	No	No
Zytiga	Oncology- Oral/Topical	Oral	Yes	No	Yes		No	Yes